

Question 1

- Describe what will happen in the nephron in response to a person drinking too much alcohol. Keep in mind that alcohol is a diuretic.

Too much alcohol?

- The first three segments of the nephron will function as usual, as well as the distal tubule. The alcohol will cause dehydration, so the collecting duct will need ADH to be produced in order to counter the diuretic alcohol. With this aquaporins are inserted in the plasma membrane, letting water pour out, therefore is more permeable to urea.

Question 2

- Describe what will happen in the nephron in response to someone who was given ecstasy and a case of bottled water.

H₂O and ecstasy

- When a person takes ecstasy the person becomes more hydrated because they want to drink more water. If there is more water, there is low blood Na⁺. The hormone released is aldosterone, which acts on the distal tubule. This activates more Na⁺ pumps and more Na⁺ is absorbed. This makes pee less concentrated.

H₂O and ecstasy

- When given ecstasy, a person will begin to consume large amounts of water. This large amount will begin to dilute the blood which causes a decrease of Na⁺. As a response aldosterone is released which acts on the distal tubule. In response to the hormone more Na⁺ pumps are activated which allows more Na⁺ to be reabsorbed...

Back to Photosynthesis!

Photosynthesis Friday!!!!




Fig. 10.11 pg. 206






Fig. 10.13 pg. 208





<http://www.youtube.com/watch?v=6BRMPrnE>




Fig. 10.13 pg 208


Use the Force

- Just like the ETC in respiration, the ETC in photosystem II proton gradient spins ATP synthase

Where does PS II happen?

If a proton gradient is created on either side of a membrane, the process must be membrane-bound

(b) Chloroplasts are highly structured, membrane-rich organelles.



Wow. Look at all of those membranes!

Copyright © 2004 Pearson Benjamin Cummings. All rights reserved.

Fig. 10.2 pg. 200


Oxygen overload!

- When Pheophytin donates an electron to the ETC, it must be replaced!
 - This drives a reaction that splits $2\text{H}_2\text{O} \rightarrow \text{O}_2 + 2\text{H}_2$
 - Oxygenic Photosynthesis
 - 2 billion years later...
 - Prokaryotic variation: $\text{H}_2\text{S} \rightarrow \text{S}_2$
 - Mini tangent:

Photosystem I

Similarities to PS II

- Both rely on the antenna complex of chlorophyll molecules



- Both transport energy via ETC...

– Iron and sulfur proteins

...until the energy reaches ferredoxin (NADP⁺ reductase)

where NADP⁺ is reduced to NADPH

Redox reactions




Fig. 10.14 pg 210

Let's Compare...

	Photosystem II	Photosystem I
Production	<input type="checkbox"/>	<input type="checkbox"/>
Wavelength	<input type="checkbox"/>	<input type="checkbox"/>

They can work alone, but work so much better together

How do they know that??

Do some Science!

Experiment

Experimental setups:

1. Expose algal cells to an intensity of far-red (700 nm) or red (680 nm) light that maximizes rate of photosynthesis at each wavelength. Then expose same algal cells to same intensities of each wavelength at same time.

Predictions: When the wavelengths are combined, the rate of photosynthesis will be double the maximum rate observed for each wavelength independently.

Prediction of null hypothesis: When the wavelengths are combined, the rate of photosynthesis will not be double the maximum rate observed for each wavelength independently.

Copyright 2004 Thomson Nelson Company. All rights reserved.

What did they measure?

Experiment

Results:

Conclusion: There is an enhancement effect for red and far-red light. The combination of 700 nm and 680 nm wavelengths more than doubles the rate of photosynthesis.

Copyright 2004 Thomson Nelson Company. All rights reserved.

Fig 10.12 pg. 207




Fig 10.15 pg. 211




Fig. 10.2 pg. 200
